



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

had Dr. DOMIN had a longer experience with our ferns, and observed the great diversity of their form and growth, he would have realized that some of his new kinds were but growths of well known species."

Nearly 70 years of experience with plants in the field, much of it with Queensland plants, add to the weight of BAILEY's comments. At the time of DOMIN's visit, the present reviewer was collecting morphological material in Queensland, and can heartily agree with the remarks regarding the variability of *Psilotum*, *Marattia*, and *Platyserium*. In *Platyserium*, particularly, the appearance of the plant is so affected by its position on the tree, that isolated plants might be described as new species, if the differences were not so obviously due, in some cases, to merely mechanical causes.—C. J. CHAMBERLAIN.

A new seed genus.—Miss BENSON²⁰ has made *Conostoma ovale* Williamson the basis of a new form genus, which she calls *Sphaerostoma*. Along with *C. ovale* the doubtful species *C. intermedium* is included as *S. ovale*. In general structure the seed resembles *Lagenostoma*, and is found most frequently without its cupule (or outer integument), from which it doubtless separated when dropped. All of the epidermal cells of the integument are papillate, and at the apex, where the integument is free from the nucellus, they become so elongated as to form a whorl of epidermal crests, the "canopy" being 8-lobed. These crests Miss BENSON calls "frills."

The structure of the lagenostome, however, is quite peculiar. There is the usual central column of nucellar tissue, surrounded by the moatlike pollen chamber which is invested by the epidermis of the nucellus. But the roof of the pollen chamber is modified into what Miss BENSON regards as "an elastically acting mechanism" which definitely closes the pollen chamber after there has been a dehiscence which admits the pollen grains. This elastic mechanism, forming the roof of the pollen chamber, is the epidermis so modified as to resemble in appearance a multiserial annulus. The conclusion is that in dehiscence this mechanism straightens elastically, admits pollen grains, and then closes the chamber again.

The seeds are provisionally referred to *Heterangium Grievii* on what seems to be excellent evidence, namely constant association, suggestions of actual continuity of ovules with the parent plant, and the evident morphological relationship to *Lagenostoma*.—J. M. C.

Culture of Opuntia.—GRIFFITHS²¹ records some very interesting observations upon the behavior of certain species of *Opuntia* under culture. The plants were grown at the government stations at Brownsville and San Antonio,

²⁰ BENSON, MARGARET J., *Sphaerostoma ovale* (*Conostoma ovale* et *intermedium* Williamson), a Lower Carboniferous ovule from Pettycur, Fifeshire, Scotland. Trans. Roy. Soc. Edinburgh 50:1-15. figs. 3. pls. 1, 2. 1914.

²¹ GRIFFITHS, D., Behavior, under cultural conditions, of species of Cacti known as *Opuntia*. U.S. Dept. Agr., Bur. Pl. Ind., Bull. no. 31. pp. 24. pls. 1-8. fig. 1. 1913.